

Third Neighbor Correlators of Spin-1/2 Heisenberg Antiferromagnet

Kazumitsu Sakai,¹ Masahiro Shiroishi,¹ Yoshihiro Nishiyama,² and Minoru Takahashi¹

¹*Institute for Solid State Physics, University of Tokyo, Kashiwa, Chiba 277-8581, Japan*

²*Department of Physics, Faculty of Science, Okayama University, Okayama 700-8530, Japan*

(Dated: February 27, 2003)

We exactly evaluate the third neighbor correlator $\langle S_j^z S_{j+3}^z \rangle$ and all the possible non-zero correlators $\langle S_j^\alpha S_{j+1}^\beta S_{j+2}^\gamma S_{j+3}^\delta \rangle$ of the spin-1/2 Heisenberg XXX antiferromagnet in the ground state without magnetic field. All the correlators are expressed in terms of certain combinations of logarithm $\ln 2$, the Riemann zeta function $\zeta(3)$, $\zeta(5)$ with rational coefficients. The results accurately coincide with the numerical ones obtained by the density-matrix renormalization group method and the numerical diagonalization.

PACS numbers: 75.10.Jm, 75.50.Ee, 02.30.Ik

Since Bethe's pioneering work of the spin-1/2 Heisenberg XXX magnet [1],

$$H = J \sum_{j=1}^L (S_j^x S_{j+1}^x + S_j^y S_{j+1}^y + S_j^z S_{j+1}^z), \quad (1)$$

exact evaluation of the correlation functions has been a long standing problem in mathematical physics. Especially significant are the spin-spin correlators $\langle S_j^z S_{j+k}^z \rangle$ (or equivalently $\langle S_j^+ S_{j+k}^- \rangle / 2$; $S_j^\pm = S_j^x \pm i S_j^y$), for which only the first and second neighbor ($k = 1, 2$) have been calculated so far:

$$\langle S_j^z S_{j+1}^z \rangle = \frac{1}{12} - \frac{1}{3} \ln 2 \simeq -0.14771572685, \quad (2)$$

$$\langle S_j^z S_{j+2}^z \rangle = \frac{1}{12} - \frac{4}{3} \ln 2 + \frac{3}{4} \zeta(3) \simeq 0.06067976996, \quad (3)$$

where $\zeta(s)$ is the Riemann zeta function and $\langle \cdots \rangle$ denotes the ground state expectation value of the antiferromagnetic model ($J > 0$). Here we have taken the thermodynamic limit $L \rightarrow \infty$. In this letter, we would like to report our new results regarding the third neighbor correlators. Our main result is

$$\begin{aligned} \langle S_j^z S_{j+3}^z \rangle &= \frac{1}{2} \langle S_j^+ S_{j+3}^- \rangle \\ &= \frac{1}{12} - 3 \ln 2 + \frac{37}{6} \zeta(3) - \frac{14}{3} \zeta(3) \ln 2 - \frac{3}{2} \zeta(3)^2 \\ &\quad - \frac{125}{24} \zeta(5) + \frac{25}{3} \zeta(5) \ln 2 \simeq -0.05024862726. \end{aligned} \quad (4)$$

In addition, we obtain the third neighbor one-particle Green function $\langle c_j^\dagger c_{j+3} \rangle_f$

$$\begin{aligned} \langle c_j^\dagger c_{j+3} \rangle_f &= \frac{1}{30} - 2 \ln 2 + \frac{169}{30} \zeta(3) - \frac{10}{3} \zeta(3) \ln 2 \\ &\quad - \frac{6}{5} \zeta(3)^2 - \frac{65}{12} \zeta(5) + \frac{20}{3} \zeta(5) \ln 2 \simeq 0.08228771669, \end{aligned} \quad (5)$$

for the isotropic spinless fermion model corresponding to (1) by the Jordan-Wigner transformation:

$$S_k^- = \prod_{j=1}^{k-1} (1 - 2c_j^\dagger c_j) c_k^\dagger, \quad S_k^+ = \prod_{j=1}^{k-1} (1 - 2c_j^\dagger c_j) c_k. \quad (6)$$

Here $\langle \cdots \rangle_f$ denotes the expectation value in the half-filled state of the spinless fermion model. Moreover we exactly calculate all the possible non-zero correlators $\langle S_j^\alpha S_{j+1}^\beta S_{j+2}^\gamma S_{j+3}^\delta \rangle$.

The result (2) comes from the ground state energy of (1) derived by Hulthén in 1938 [2]. The result (3) was obtained by one of the authors in 1977 [3, 4] via the strong coupling expansion for the ground state energy of the half-filled Hubbard model. This result is also reproduced in the framework of the asymptotic Bethe ansatz for an integrable spin chain with variable range exchange [5]. However, probably due to the complexity of the wave function for these models, no one has succeeded in generalizing the method to obtain the higher neighbor correlators.

On the other hand, utilizing the representation theory of the quantum affine algebra $U_q(\widehat{sl_2})$ and the associated vertex operators, in 1992, Jimbo *et al.* derived a universal multiple integral representation of arbitrary correlators for the massive XXZ antiferromagnet [6, 7]. Their result has been extended to the XXX [8, 9], the massless XXZ [10, 11] and the XYZ [12] antiferromagnets. However the explicit evaluation, even for the second neighbor correlator (3), was not achieved for a long time.

In this respect, it is quite remarkable that Boos and Korepin recently devised a general method to evaluate the multiple integral representation especially in the study of the Emptiness Formation Probability (EFP) for the XXX antiferromagnet [13, 14]. The EFP, $P(n)$ describes the probability of finding a ferromagnetic string of length n in the antiferromagnetic ground state [9]. Explicitly it reads

$$P(n) = \left\langle \prod_{j=1}^n \left(S_j^z + \frac{1}{2} \right) \right\rangle. \quad (7)$$

By reducing the integrand of the multiple integral representation to certain *canonical form*, the EFP for $n = 3, 4$ [13, 14] and $n = 5$ [15] were evaluated by Boos *et al.* (see also recent progress for $n = 6$ [16]). Note that $P(2)$ and $P(3)$ in (7) are related to the first and sec-

and neighbor correlators as $P(2) = 1/4 + \langle S_j^z S_{j+1}^z \rangle$ and $P(3) = 1/8 + \langle S_j^z S_{j+1}^z \rangle + \langle S_j^z S_{j+2}^z \rangle / 2$.

Here we quote the explicit form of $P(4)$ obtained in [13, 14], which is closely related to the third neighbor correlator $\langle S_j^z S_{j+3}^z \rangle$.

$$\begin{aligned} P(4) &= \frac{1}{16} + \frac{3}{4} \langle S_j^z S_{j+1}^z \rangle + \frac{1}{2} \langle S_j^z S_{j+2}^z \rangle + \frac{1}{4} \langle S_j^z S_{j+3}^z \rangle \\ &\quad + \langle S_j^z S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle \\ &= \frac{1}{5} - 2 \ln 2 + \frac{173}{60} \zeta(3) - \frac{11}{6} \zeta(3) \ln 2 - \frac{51}{80} \zeta(3)^2 \\ &\quad - \frac{55}{24} \zeta(5) + \frac{85}{24} \zeta(5) \ln 2. \end{aligned} \quad (8)$$

Note that on the antiferromagnetic ground state without magnetic field, all the correlators with an odd number of S^z vanishes. Substituting (2) and (3) into (8), one finds the relation between the third neighbor correlator $\langle S_j^z S_{j+3}^z \rangle$ and the four point correlator $\langle S_j^z S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle$. However the exact value of $\langle S_j^z S_{j+3}^z \rangle$ itself can not be determined solely from $P(4)$.

To determine $\langle S_j^z S_{j+3}^z \rangle$, we consider the following auxiliary correlator:

$$\begin{aligned} P_{+-+-}^{+-} &= \frac{1}{16} - \frac{3}{4} \langle S_j^z S_{j+1}^z \rangle + \frac{1}{2} \langle S_j^z S_{j+2}^z \rangle - \frac{1}{4} \langle S_j^z S_{j+3}^z \rangle \\ &\quad + \langle S_j^z S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle. \end{aligned} \quad (9)$$

Here and hereafter $P_{\varepsilon_1 \varepsilon_2 \varepsilon_3 \varepsilon_4}^{\tilde{\varepsilon}_1 \tilde{\varepsilon}_2 \tilde{\varepsilon}_3 \tilde{\varepsilon}_4}$ (also written as $P_{\varepsilon}^{\tilde{\varepsilon}}$ for simplicity) denotes a correlator of the form

$$P_{\varepsilon_1 \varepsilon_2 \varepsilon_3 \varepsilon_4}^{\tilde{\varepsilon}_1 \tilde{\varepsilon}_2 \tilde{\varepsilon}_3 \tilde{\varepsilon}_4} = \langle E_1^{\tilde{\varepsilon}_1 \varepsilon_1} E_2^{\tilde{\varepsilon}_2 \varepsilon_2} E_3^{\tilde{\varepsilon}_3 \varepsilon_3} E_4^{\tilde{\varepsilon}_4 \varepsilon_4} \rangle, \quad (10)$$

where $\varepsilon_j, \tilde{\varepsilon}_j = \{+, -\}$ and $E_j^{\varepsilon_j \tilde{\varepsilon}_j}$ is the 2×2 elementary matrix ($E^{\pm\pm} = \pm S^z + 1/2$, $E^{-+} = S^+$, $E^{+-} = S^-$) acting on j th site. In this notation $P(4) = P_{++++}^{++++}$. Note that because the Hamiltonian (1) has the symmetry under the group $SU(2)$, $P_{\varepsilon}^{\tilde{\varepsilon}}$ possesses a property like

$$P_{\varepsilon}^{\tilde{\varepsilon}} = P_{\tilde{\varepsilon}}^{\varepsilon} = P_{-\varepsilon}^{-\tilde{\varepsilon}} = P_{-\tilde{\varepsilon}}^{-\varepsilon}. \quad (11)$$

As in the case of $P(4)$, the correlators (10) enjoy the multiple integral representation [6, 7, 8, 10]:

$$P_{\varepsilon}^{\tilde{\varepsilon}} = \prod_{j=1}^4 \int_C \frac{d\lambda_j}{2\pi i} U(\lambda_1, \dots, \lambda_4) T(\lambda_1, \dots, \lambda_4), \quad (12)$$

where the integration contour C is taken to be a line $[-\infty - i\alpha, \infty - i\alpha]$ ($0 < \alpha < 1$). For convenience, we choose $\alpha = 1/2$. The integrand $U(\lambda_1, \dots, \lambda_4)$ is given by

$$U(\lambda_1, \dots, \lambda_4) = \pi^{10} \frac{\prod_{1 \leq k < j \leq 4} \sinh \pi \lambda_{jk}}{\prod_{j=1}^4 \sinh^4 \pi \lambda_j}, \quad (13)$$

while $T(\lambda_1, \dots, \lambda_4)$ depends on the selection of ε and $\tilde{\varepsilon}$. Here and hereafter we use the notation $\lambda_{jk} = \lambda_j -$

λ_k to save space. In particular for the correlator (9), $T(\lambda_1, \dots, \lambda_4)$ is given by

$$T(\lambda_1, \dots, \lambda_4) = \frac{\lambda_1(\lambda_1 + i)^2 \lambda_2^3 \lambda_3(\lambda_3 + i)^2 \lambda_4^3}{(\lambda_{21} - i) \lambda_{31} \lambda_{41} \lambda_{32} \lambda_{42} (\lambda_{43} - i)}. \quad (14)$$

To calculate the multiple integral (12), we follow the method by Boos and Korepin [14]. Roughly, their method is described as follows. First Taking carefully into account the property of $U(\lambda_1, \dots, \lambda_4)$, we modify the integrand $T(\lambda_1, \dots, \lambda_4)$ such that the integral gives the same result as the original one ("weak equivalence"). In this way it is likely that the integrand $T(\lambda_1, \dots, \lambda_4)$ can always be reduced to the following form (we call it "canonical form"):

$$T_c = P_0(\lambda_2, \lambda_3, \lambda_4) + \frac{P_1(\lambda_1, \lambda_3, \lambda_4)}{\lambda_{21}} + \frac{P_2(\lambda_1, \lambda_3)}{\lambda_{21} \lambda_{43}}, \quad (15)$$

where P_0 , P_1 and P_2 are certain polynomials. Once one derives the canonical form, one can perform the multiple integral (12) by using the Cauchy theorem [14]. The result is written as combinations of the logarithm $\ln 2$, the Riemann zeta function $\zeta(3)$ and $\zeta(5)$ and rational numbers. Consequently, the main part of the calculation for the multiple integral (12) reduces to finding the canonical form (15).

Now we consider the case (14) and show that the Boos-Korepin method is also applicable to our case. Let us introduce the following diagram.

$$j \xrightarrow[k]{k} := \frac{1}{\lambda_{jk}}, \quad j \xrightarrow[k]{k} := \frac{1}{\lambda_{jk} - i},$$

where $j > k$. First we expand (14) through partial fractions. The result consists of 24 terms. Taking into account the antisymmetry of the function (13) under transposition of any two variables λ_j and λ_k and the symmetry of T (14) under $\lambda_1 \leftrightarrow \lambda_3$, $\lambda_2 \leftrightarrow \lambda_4$, we can reduce the 24 partial fractions to eight ones. Diagrammatically, its denominators are written as

$$\begin{aligned} & \begin{array}{c} 1 \quad 2 \\ \diagdown \quad \diagup \\ 4 \quad 3 \end{array} \rightarrow \begin{array}{c} 1 \quad 2 \\ \diagup \quad \diagdown \\ 4 \quad 3 \end{array} - \begin{array}{c} 1 \quad 2 \\ \diagdown \quad \diagup \\ 4 \quad 3 \end{array} + i \left(\begin{array}{c} 1 \quad 2 \\ \diagup \quad \diagup \\ 4 \quad 3 \end{array} + \begin{array}{c} 1 \quad 2 \\ \diagdown \quad \diagdown \\ 4 \quad 3 \end{array} \right) \\ & + 2i \left(\begin{array}{c} 1 \quad 2 \\ \diagup \quad \diagup \\ 4 \quad 3 \end{array} - \begin{array}{c} 1 \quad 2 \\ \diagdown \quad \diagdown \\ 4 \quad 3 \end{array} - \begin{array}{c} 1 \quad 2 \\ \diagup \quad \diagdown \\ 4 \quad 3 \end{array} - \begin{array}{c} 1 \quad 2 \\ \diagdown \quad \diagup \\ 4 \quad 3 \end{array} \right). \end{aligned}$$

From a symmetry of the denominator, the second term can further be simplified as

$$\begin{aligned} & -g \begin{array}{c} 1 \quad 2 \\ \diagdown \quad \diagup \\ 4 \quad 3 \end{array} \rightarrow -\frac{1}{4} (g(\lambda_1, \lambda_3, \lambda_2, \lambda_4) - g(\lambda_4, \lambda_1, \lambda_3, \lambda_2) \\ & + g(\lambda_2, \lambda_4, \lambda_1, \lambda_3) - g(\lambda_3, \lambda_2, \lambda_4, \lambda_1)) \begin{array}{c} 1 \quad 2 \\ \diagup \quad \diagup \\ 4 \quad 3 \end{array} \\ & \rightarrow \frac{1}{2} \lambda_1 \lambda_2 \lambda_3 \lambda_4 (i\lambda_1 + i\lambda_3 + 2\lambda_1 \lambda_3) (i\lambda_2 + i\lambda_4 + 2\lambda_2 \lambda_4) \\ & \times 1 \xrightarrow[k]{k} 2 \times 4 \xrightarrow[k]{k} 3, \end{aligned}$$

where g denotes the numerator of (14). Next using the Cauchy theorem, we shift variables $\lambda_j \rightarrow \lambda_j \pm i$ such that the denominators do not contain i . For instance, we have

$$\begin{aligned} g &= g^{(1)} + g^{(2)} \\ &\rightarrow g^{(1)} + g_{\lambda_2 \rightarrow \lambda_2 + i}^{(2)}, \end{aligned}$$

where

$$\begin{aligned} g^{(1)} &= -\lambda_1(\lambda_1 + i)\lambda_2^3\lambda_3(\lambda_3 + i)^2\lambda_4^3 \\ g^{(2)} &= -\lambda_1(\lambda_1 + i)\lambda_2^4\lambda_3(\lambda_3 + i)^2\lambda_4^3. \end{aligned}$$

Finally again using the antisymmetric property of (13), we eliminate the symmetric part with respect to $\lambda_j \leftrightarrow \lambda_k$. In this way we have obtain the canonical form as

$$\begin{aligned} P_0 &= \frac{56}{5}\lambda_2\lambda_3^2\lambda_4^3, \\ P_1 &= \frac{27}{10}\lambda_4 - i\lambda_4^2 + \frac{33}{5}\lambda_3\lambda_4^2 + \frac{4}{5}\lambda_4^3 + 2i\lambda_3\lambda_4^3 + 4\lambda_3^2\lambda_4^3 \\ &\quad + \lambda_1(-4i\lambda_4 + 7\lambda_4^2 - 32i\lambda_3\lambda_4^2 - 10i\lambda_4^3 - 12i\lambda_3^2\lambda_4^3) \\ &\quad + \lambda_1^2(4\lambda_4 - 19i\lambda_4^2 - 28\lambda_3\lambda_4^2 - 10\lambda_4^3 \\ &\quad \quad - 28i\lambda_3\lambda_4^3 - 32\lambda_3^2\lambda_4^3), \\ P_2 &= -\frac{3}{10} + \frac{3}{2}i\lambda_3 + \frac{3}{2}\lambda_1\lambda_3 - \frac{1}{2}\lambda_3^2 + 4i\lambda_1\lambda_3^2 + 6\lambda_1^2\lambda_3^2. \end{aligned}$$

Subsequently, applying the method developed in [14], we calculate the multiple integral by substituting the above canonical form into (12). Explicitly,

$$\begin{aligned} P_{+-+}^{+-+} &= J_0 + J_1 + J_2 \\ J_0 &= \frac{7}{10}, \quad J_1 = -\frac{2}{3} + \frac{3}{10}\zeta(3) + \frac{35}{32}\zeta(5), \\ J_2 &= -\frac{1}{2}\zeta(3) + \frac{1}{2}\zeta(3)\ln 2 + \frac{9}{80}\zeta(3)^2 - \frac{25}{32}\zeta(5) \\ &\quad - \frac{5}{8}\zeta(5)\ln 2, \end{aligned} \quad (16)$$

where J_k denotes the result of the integration regarding the term P_k . We remark that the canonical form is not unique due to the non-uniqueness of partial fraction expansions. Accordingly, the explicit value of each J_k depends on the choice of the canonical form. The final result $J_0 + J_1 + J_2$, however, is always unique as a matter of course.

Combining the result (16) with (8) and (9), we obtain the third neighbor correlator $\langle S_j^z S_{j+3}^z \rangle$ (4) and at the same time the correlator $\langle S_j^z S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle$ as

$$\begin{aligned} \langle S_j^z S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle &= \frac{1}{80} - \frac{1}{3}\ln 2 + \frac{29}{30}\zeta(3) - \frac{2}{3}\zeta(3)\ln 2 \\ &\quad - \frac{21}{80}\zeta(3)^2 - \frac{95}{96}\zeta(5) + \frac{35}{24}\zeta(5)\ln 2. \end{aligned} \quad (17)$$

Now let us consider the four-point correlators of the form $\langle S_j^\alpha S_{j+1}^\beta S_{j+2}^\gamma S_{j+3}^\delta \rangle$ ($\{\alpha, \beta, \gamma, \delta\} \in \{x, y, z, 0\}$; $S_j^0 = \mathbb{1}$). Because the correlators with an odd number of $S^{\{\alpha\}} \setminus 0$ vanish, the possible non-zero correlators are restricted to the following three types: $\langle S_j^\alpha S_{j+1}^\alpha S_{j+2}^\beta S_{j+3}^\beta \rangle$, $\langle S_j^\alpha S_{j+1}^\beta S_{j+2}^\alpha S_{j+3}^\beta \rangle$ and $\langle S_j^\alpha S_{j+1}^\beta S_{j+2}^\beta S_{j+3}^\alpha \rangle$. Further due to the isotropy of the Hamiltonian (1), one can find that the independent correlators are written as the following seven ones: $\langle S_j^x S_{j+1}^x S_{j+2}^x S_{j+3}^x \rangle$, $\langle S_j^z S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle$, $\langle S_j^z S_{j+1}^x S_{j+2}^x S_{j+3}^z \rangle$ and already obtained ones in (2)–(4) and (17). Then we shall calculate the remaining three correlators here. For convenience we use the operator $S^\pm (= S^x \pm iS^y)$ instead of S^x . First we consider the correlator $\langle S_j^+ S_{j+1}^z S_{j+2}^z S_{j+3}^- \rangle (= 2\langle S_j^z S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle)$. From (10) and the property (11), this correlator is expressed as $\langle S_j^+ S_{j+1}^z S_{j+2}^z S_{j+3}^- \rangle = (P_{-+++}^{++++} - P_{-+-+}^{++++})/2$. Using the relation $\langle S_j^+ S_{j+3}^- \rangle = 2(P_{-+++}^{++++} + P_{-+-+}^{++++})$, we obtain

$$\langle S_j^+ S_{j+1}^z S_{j+2}^z S_{j+3}^- \rangle = P_{-+++}^{++++} - \frac{1}{2}\langle S_j^z S_{j+3}^z \rangle. \quad (18)$$

Here we have used the relation $\langle S_j^+ S_{j+3}^- \rangle = 2\langle S_j^z S_{j+3}^z \rangle$. Similarly the other correlators are given by

$$\begin{aligned} \langle S_j^z S_{j+1}^+ S_{j+2}^z S_{j+3}^- \rangle &= P_{-+++}^{++++} - \frac{1}{2}\langle S_j^z S_{j+2}^z \rangle, \\ \langle S_j^z S_{j+1}^z S_{j+2}^+ S_{j+3}^- \rangle &= P_{-+++}^{++++} - \frac{1}{2}\langle S_j^z S_{j+1}^z \rangle. \end{aligned} \quad (19)$$

Therefore our goal is to evaluate the auxiliary correlators P_{-+++}^{++++} , P_{-+-+}^{++++} and P_{-+-+}^{++++} . They are given if we replace the integrand $T(\lambda_1, \dots, \lambda_4)$ by

$$T^{(l)} = \frac{(\lambda_1 + i)^3 \lambda_2 (\lambda_2 + i)^2 (\lambda_3 + i) \lambda_3^2 \lambda_4^{4-l} (\lambda_4 + i)^{l-1}}{(\lambda_{21} - i)(\lambda_{31} - i)(\lambda_{32} - i) \lambda_{41} \lambda_{42} \lambda_{43}}.$$

in the multiple integral representation. Here the correlator P_{-+++}^{++++} , P_{-+-+}^{++++} and P_{-+-+}^{++++} correspond to $l = 1, 2$ and 3, respectively. Using the procedure similar to the case of P_{-+-+}^{++++} , one obtains the explicit values of the above auxiliary correlators. As a result, combining the identity (18) and (19) with (2)–(4), we arrive at

$$\begin{aligned} \langle S_j^+ S_{j+1}^z S_{j+2}^z S_{j+3}^- \rangle &= \frac{1}{120} - \frac{1}{2}\ln 2 + \frac{169}{120}\zeta(3) \\ &\quad - \frac{5}{6}\zeta(3)\ln 2 - \frac{3}{10}\zeta(3)^2 - \frac{65}{48}\zeta(5) + \frac{5}{3}\zeta(5)\ln 2, \end{aligned} \quad (20)$$

$$\begin{aligned} \langle S_j^+ S_{j+1}^z S_{j+2}^- S_{j+3}^z \rangle &= \frac{1}{120} - \frac{1}{3}\ln 2 + \frac{77}{60}\zeta(3) \\ &\quad - \frac{5}{6}\zeta(3)\ln 2 - \frac{3}{10}\zeta(3)^2 - \frac{65}{48}\zeta(5) + \frac{5}{3}\zeta(5)\ln 2, \end{aligned} \quad (21)$$

$$\begin{aligned} \langle S_j^+ S_{j+1}^- S_{j+2}^z S_{j+3}^z \rangle &= \frac{1}{120} + \frac{1}{6}\ln 2 - \frac{91}{120}\zeta(3) \\ &\quad + \frac{1}{3}\zeta(3)\ln 2 + \frac{3}{40}\zeta(3)^2 + \frac{35}{48}\zeta(5) - \frac{5}{12}\zeta(5)\ln 2. \end{aligned} \quad (22)$$

We mention a few remarks of our results. (i) All the above correlators are written as the logarithm $\ln 2$, the

TABLE I: Estimates of the correlators by the exact evaluations, DMRG and the extrapolations from the numerical diagonalization for the system size $L = 24, 28, 32$.

Correlators	Exact	DMRG	Extrap.
$\langle S_j^z S_{j+3}^z \rangle$	-0.0502486	-0.0502426	-0.0502475
$\langle S_j^+ S_{j+1}^z S_{j+2}^z S_{j+3}^- \rangle^a$	0.0205719	0.0205681	0.0205716
$\langle S_j^z S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle$	0.0307153	0.0307105	0.0307154
$\langle S_j^+ S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle$	-0.0141607	-0.0141579	-0.0141606
$\langle S_j^+ S_{j+1}^z S_{j+2}^z S_{j+3}^z \rangle$	0.0550194	0.0550108	0.0550198
$^a \frac{1}{4} \langle c_j^\dagger c_{j+3} \rangle_f$			

Riemann zeta function $\zeta(3)$ and $\zeta(5)$. This agrees with the general conjecture by Boos and Korepin: *arbitrary correlators of the XXX antiferromagnet are described as certain combinations of logarithm $\ln 2$, the Riemann zeta function with odd arguments and rational coefficients*. Especially intriguing is the existence of the non-linear terms such as $\zeta(3)^2$, $\zeta(3) \ln 2$ and $\zeta(5) \ln 2$. (ii) The correlator (20) is interpreted as the third neighbor one-particle Green functions $\langle c_j^\dagger c_{j+3} \rangle_f / 4$ via the Jordan-Wigner transformation (6). Obviously, the first neighbor one-particle Green function is expressed as

$$\langle c_j^\dagger c_{j+1} \rangle_f = \frac{1}{6} - \frac{2}{3} \ln 2 \simeq -0.295431453707,$$

which coincides with $\langle S_j^+ S_{j+1}^- \rangle$. Due to the characteristic $\pi/2$ -oscillation: $\langle c_j^\dagger c_k \rangle_f \sim A_{jk} \cos(\pi(k-j+1)/2)$, one finds $\langle c_j^\dagger c_{j+2k} \rangle_f = 0$. Therefore the quantity (5) is the first non-trivial exact result of the correlators containing the fermionic nature. (iii) The difference between (20) and (21) gives the nearest chiral correlator

$$\begin{aligned} \langle (\mathbf{S}_j \times \mathbf{S}_{j+1}) \cdot (\mathbf{S}_{j+2} \times \mathbf{S}_{j+3}) \rangle &= 3 \langle (S_j^+ S_{j+1}^z S_{j+2}^- S_{j+3}^z) \rangle \\ &- \langle S_j^+ S_{j+1}^z S_{j+2}^z S_{j+3}^- \rangle = \frac{1}{2} \ln 2 - \frac{3}{8} \zeta(3), \end{aligned}$$

which exactly agrees with the one derived from the ground state energy of an integrable two-chain model with four-body interactions [17].

To confirm the validity of our formulae, we performed numerical calculations by using the density-matrix renormalization group (DMRG) [18, 19] and numerical diagonalization. As for the DMRG, we followed standard algorithm [20]. We have repeated renormalization 500-times. At each renormalization, we kept, at most, 200 relevant states for a (new) block. The numerical diagonalization was performed for the system size $L = 24, 28$ and 32. We extrapolate the data from a fitting function $a_0 + a_1/L^2 + a_2/L^4$. All our analytical results coincide quite accurately with both numerical ones (TABLE I).

In closing we would like to comment on generalizations of the present results. The extension to the calculation of higher neighbor correlators $\langle S_j^z S_{j+k}^z \rangle_{k \geq 4}$ is of great interest. The fourth neighbor one $\langle S_j^z S_{j+4}^z \rangle$, for example, will

be calculated by combination of the EFP, $P(5)$ and two independent auxiliary correlators, which can in principle be evaluated. In fact $P(5)$ has been already obtained in [15]. The computation, however, will be much more complicated. Alternatively, extending the present result to the inhomogeneous case as in [16] and taking into account the property of the quantum Knizhnik-Zamolodchikov equation, we may derive higher neighbor correlators. Using this, eventually we hope to extract the long-distance asymptotics $\langle S_j^z S_{j+k}^z \rangle_{k \gg 1}$, which is a crucial problem in conformal field theory [21, 22].

The authors are grateful to H.E. Boos and V.E. Korepin for many valuable discussions. KS is supported by the JSPS research fellowships for young scientists. MS and YN are supported by Grant-in-Aid for young scientists No. 14740228 and No. 13740240, respectively. This work is in part supported by Grant-in Aid for the Scientific Research (B) No. 14340099 from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

-
- [1] H.A. Bethe, Z. Phys. **71**, 205 (1931).
 - [2] L. Hulthén, Ark. Mat. Astron. Fys. A **26**, 1 (1938).
 - [3] M. Takahashi, J. Phys. C **10**, 1289 (1977).
 - [4] M. Takahashi, *Thermodynamics of One-Dimensional Solvable Models*, (Cambridge University Press, Cambridge, 1999).
 - [5] J. Dittrich and V.I. Inozemtsev, J. Phys. A **30**, L623 (1997).
 - [6] M. Jimbo, K. Miki, T. Miwa, and, A. Nakayashiki, Phys. Lett. A **168**, 256 (1992).
 - [7] M. Jimbo and T. Miwa, *Algebraic Analysis of Solvable Lattice Models*, (American Mathematical Society, Providence, RI, 1995).
 - [8] A. Nakayashiki, Int. J. Mod. Phys. A **9**, 5673 (1994).
 - [9] V.E. Korepin, A.G. Izergin, F.H.L. Eßler, and D.B. Uglov, Phys. Lett. A **190**, 182 (1994).
 - [10] M. Jimbo and T. Miwa, J. Phys. A **29**, 2923 (1996).
 - [11] N. Kitanine, J.M. Maillet, and V. Terras, Nucl. Phys. B **567**, 554 (2000).
 - [12] Y-H. Quano, J. Phys. A **35**, 9549 (2002).
 - [13] H.E. Boos and V.E. Korepin, J. Phys. A **34**, 5311 (2001).
 - [14] H.E. Boos and V.E. Korepin, *Integrable models and Beyond*, edited by M. Kashiwara and T. Miwa (Birkhäuser, Boston, 2002); hep-th/0105144.
 - [15] H.E. Boos, V.E. Korepin, Y. Nishiyama, and M. Shiroyishi, J. Phys. A **35** 4443 (2002).
 - [16] H.E. Boos, V.E. Korepin, and F.A. Smirnov, hep-th/0209246.
 - [17] N. Muramoto and M. Takahashi, J. Phys. Soc. Jpn. **68**, 2098 (1999).
 - [18] S. R. White, Phys. Rev. Lett. **69**, 2963 (1992).
 - [19] S. R. White, Phys. Rev. B **48**, 10345 (1993).
 - [20] *Density-Matrix Renormalization: A New Numerical Method in Physics*, edited by I. Peschel, X. Wang, M. Kaulke and K. Hallberg (Springer-Verlag, Berlin, 1999).
 - [21] I. Affleck, J. Phys. A **31**, 4573 (1998).
 - [22] S. Lukyanov and V. Terras, hep-th/0206093.